#### International Workshop on NEW PARTIAL WAVE ANALYSIS TOOLS FOR NEXT GENERATION HADRON SPECTROSCOPY EXPERIMENTS June 20-22, 2012 Camogli, Italy

#### PWA approach for j/psi decays at BES

-----Automatic Generation of the Partial Wave Analysis Program For Physics Processes

Jian-Xiong Wang Institute of High Energy Physics, Academia Sinica, Beijing, China email: jxwang@ihep.ac.cn

- 1. Brief Introduction to FDC
- 2. The Rules to Construct the Effective Lagrangian
- 3. FDC-PWA application
- 4. Summary

1. Brief Introduction

Feynman Diagram Calculation(FDC), The project was started from 1993.

PROGRESS IN FDC PROJECT. Nucl.Instrum.Meth.A534:241-245,2004

FDC Homepage: http://v-www.ihep.ac.cn/~wjx

FDC-SM-and-Many-Extensions FDC-NRQCD FDC-MSSM FDC-PWA FDC-LOOP There is a common kernel. The Source is written in REDUCE, RLISP.

FDC-PWA was started from 1998 and constructed for BES partial wave analysis. PWA: to decide the  $J^{PC}$ , mass, width and branch ratio of a resonant.

#### FDC-PWA: Powerful Tool

To work with high spin states (0, 1/2, 1, 3/2, 2, 5/2, 3, 7/2, 4, 9/2) and construct effective Lagrangians.

The expression of the effective interaction vertices and the propagators for the high spin states are quite lengthy.

The related amplitudes and amplitude squares are complicated.

There are many free parameters in the effective Lagrangian and these parameters will be fixed when the generated program is used to do Likelihood fitting of experimental data.

To generate a complete set of the Fortran sources to do the partial wave analysis on experimental data.

## 2. The Rule to Construct Effective Lagrangian For PWA



For NN scatting, T-channel vertices must be considered:



The Coefficients of independent terms depend on the:  $t=(PA - PB)^2$  Option to apply Isopin symmetry:





The propagator for resonance:

$$\frac{1}{p^2 - m^2 + im\Gamma}$$

Suppersion Factor for interaction vertices when the resonance is off mass shell:

Barrier, 
$$-\left|p^2 - m^2\right|/a$$

There is an option to choose the suppersion factor

# **Physical Process**

Input for a physical process: process, physical model, Many options, histograms, scatter plots.

Generate Feynman Diagram

Manipulate amplitudes for each diagram and generate FORTRAN source for calculation of amplitudes square

Find and properly treat all the resonnance, t-channel

singularities, ... and generate FORTRAN source for phase space integral and

fitting

Control flag and parameters files generated by FDC which can be changed later by users: flag.inp, amptable.inp, fpara.inp, reson.inp

Users should prepare two files: pdata1.dat –

experiment events data file pdata1.mc -

phase space monte carlo event file

Compile FORTRAN programs and run 'fit' for likelihood fitting

Output: mplot.info, pep.res, mplot.hbook, dplot.hbook

## 3. FDC-PWA Application:

- . > gmodel
- . > doall
- . > cd fort
- . > make
- . > fit

FDC-PWA: http://v-www.ihep.ac.cn/~wjx/pwa/index.html

FDC-PWA Application For BEPC physics:

Generated a Effective Lagrangian at BES energy range (J/psi) with about 300 resonant (including high spin state up to spin=7/2)

Generated all the 2, 3, 4, 5, 6 final state decays of J/psi.

Used to do PWA analysis by BES Experimental Group.

Event Generator: There are event generator for each generated physicalprocess, with the free parameters wating for PWA to fix.



## 4. Summary

We supply a powerful tool: FDC-PWA http://v-www.ihep.ac.cn/~wjx/pwa/index.html

- . To constructe the effective Lagrangian.
- To generate the complete set of Fortran source for PWA analysis.
- . To generate an event generator.
- . be used in BES PWA analysis since 1999.
- . Could be used for any possible experiment.

Thanks!

## **FDC-PWA Application on CSR Physics:**



Example:

p+p---->p+p

p+p---->p+n+pi

p+p---->p+p+pi

### p+p--->p+p







O of proton and beam in lab

## p+p--->p+n+pi

P

 $\mathbf{P}$ 

=

π.

o[770]

 $\mathbf{P}$ 

 $\mathbf{P}$ 

Diagram 23

 $\mathbf{p}$ 

p.





Diagram 24



P

Diagram 10

 $\Delta[1620]_{+1}^{n}$ 



Diagram 15



Diagram 20







O of n and beam in lab

O of p and beam in lab

O of pi and beam in lab

### p+p--->p+p+pi









