

The Gamma function

Marco Bonvini

October 9, 2010

1 Gamma function

The Euler Gamma function is defined as

$$\Gamma(z) \equiv \int_0^{\infty} dt e^{-t} t^{z-1} . \quad (1)$$

It is easy to show that $\Gamma(z)$ satisfy the recursion relation

$$\Gamma(z + 1) = z \Gamma(z) : \quad (2)$$

indeed, integrating by parts,

$$\begin{aligned} \Gamma(z + 1) &= \int_0^{\infty} dt e^{-t} t^z \\ &= -e^{-t} t^z \Big|_0^{\infty} + z \int_0^{\infty} dt e^{-t} t^{z-1} \\ &= z \Gamma(z) . \end{aligned} \quad (3)$$

For $z = n$ integer eq. (2) is the recursion relation of the factorial, and thus we have

$$\Gamma(n + 1) \propto n! ; \quad (4)$$

because in addition $\Gamma(1) = 1$ (easily derived from the definition), we have the identification

$$\Gamma(n + 1) = n! , \quad (5)$$

and in this sense the Gamma function is a complex extension of the factorial.

The sequence of Gamma function computed in all half-integers can be obtained using subsequently the recursion relation (2) and knowing that

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi} \quad (6)$$

that is easy to compute:

$$\begin{aligned} \Gamma\left(\frac{1}{2}\right) &= \int_0^{\infty} dt e^{-t} t^{-\frac{1}{2}} \\ &= 2 \int_0^{\infty} dx e^{-x^2} \\ &= 2 \frac{\sqrt{\pi}}{2} = \sqrt{\pi} . \end{aligned} \quad (7)$$




Figure 1: $\text{Re } \Gamma(x)$ for real x .

1.1 Analytical structure

First, from the definition (1), we see that

$$\boxed{\Gamma(\bar{z}) = \overline{\Gamma(z)}} , \quad (8)$$

that is to say that Gamma is a real function, and in particular $\text{Im } \Gamma(x) = 0$ for $x \in \mathbb{R}$.

Inverting eq. (2) we have

$$\Gamma(z) = \frac{1}{z} \Gamma(z+1) , \quad (9)$$

and when $z = 0$ it diverges (because $\Gamma(1) = 1$ is finite). It is evident that $z = 0$ corresponds to a simple pole of order 1 and residue 1: indeed

$$\text{Res } \Gamma(z) = \lim_{z \rightarrow 0} z \Gamma(z) = \lim_{z \rightarrow 0} \Gamma(z+1) = 1 . \quad (10)$$

Formally, we can discover that the Gamma function has simple poles in all negative integers simply iterating the recursion (2):

$$\begin{aligned} \Gamma(-n) &= \frac{1}{-n} \Gamma(-n+1) \\ &= \dots \\ &= \frac{(-1)^n}{n!} \Gamma(0) \quad (n \in \mathbb{N}) . \end{aligned} \quad (11)$$

This brings to the conclusion that $\Gamma(-n)$ has a simple pole of order 1 and residue $\frac{(-1)^n}{n!}$. More rigorously, this can be shown in the usual way:

$$\begin{aligned} \text{Res } \Gamma(z-n) &= \lim_{z \rightarrow 0} z \Gamma(z-n) \\ &= \lim_{z \rightarrow 0} \frac{z}{z-n} \Gamma(z-n+1) \\ &= \dots \end{aligned}$$

$$\begin{aligned}
&= \lim_{z \rightarrow 0} \frac{z}{(z-n)(z-n+1)\cdots(z-1)} \Gamma(z) \\
&= \frac{(-1)^n}{n!} \quad (n \in \mathbb{N}).
\end{aligned} \tag{12}$$

In all the complex plane (except the negative real axis) the Gamma function is well defined. For non-integer negative real values the Gamma function can be analytically continued (as we have seen for example for half-integers, positives and negatives).

We can prove these results in a simpler way. Starting from the definition (1) we rewrite

$$\begin{aligned}
\Gamma(z) &= \int_1^\infty dt e^{-t} t^{z-1} + \int_0^1 dt e^{-t} t^{z-1} \\
&= \int_1^\infty dt e^{-t} t^{z-1} + \int_0^1 dt t^{z-1} \sum_{n=0}^\infty \frac{(-1)^n}{n!} t^n \\
&= \int_1^\infty dt e^{-t} t^{z-1} + \sum_{n=0}^\infty \frac{(-1)^n}{n!} \frac{1}{z+n}
\end{aligned} \tag{13}$$

where the first integral is an analytic function of z and the second term gives the position of the poles and their residues.

Moreover, the Gamma function has an essential singularity to complex infinity, because $\Gamma\left(\frac{1}{z}\right)$ has a non-defined limit for $z \rightarrow 0$. This is to say that the Gamma function is not well defined in the compactified complex plane.

Now we want to show that near a pole in $-n$ one has the expansion

$$\boxed{\Gamma(z-n) = \frac{(-1)^n}{n!} \left[\frac{1}{z} + \psi(n+1) + \mathcal{O}(z) \right]} \tag{14}$$

where $\psi(z)$ is the logarithmic derivative of the Gamma function, defined in (41); note that $\psi(n+1)$ has an explicit expression given in eq. (44). First, note that

$$\begin{aligned}
\Gamma(z-n) &= \frac{1}{z-n} \Gamma(z-n+1) \\
&= \dots \\
&= \frac{1}{(z-n)(z-n+1)\cdots(z-1)z(z+1)\cdots(z+n)} \Gamma(z+n+1) \\
&= \frac{1}{z} G(z,n)
\end{aligned} \tag{15}$$

where we have defined

$$G(z,n) = \frac{\Gamma(z+n+1)}{(z^2-n^2)\cdots(z^2-1)} = \Gamma(z+n+1) \frac{(-1)^n}{(n!)^2} \left[1 + \mathcal{O}(z^2) \right]. \tag{16}$$

By Taylor expanding $G(z,n)$ around $z=0$ in (15) we get

$$\begin{aligned}
\Gamma(z-n) &= \frac{1}{z} \left[G(0,n) + z G'(0,n) + \mathcal{O}(z^2) \right] \\
&= \frac{(-1)^n}{n!} \left[\frac{1}{z} + \frac{1}{n!} \Gamma'(n+1) + \mathcal{O}(z) \right],
\end{aligned} \tag{17}$$

that is the claimed result.

1.2 Alternative definitions

There are some alternative definitions of the Gamma function. One, due to Euler, is

$$\Gamma(z) = \frac{1}{z} \prod_{n=1}^{\infty} \frac{(1 + \frac{1}{n})^z}{1 + \frac{z}{n}} = \lim_{n \rightarrow \infty} \frac{n! (n+1)^z}{z(z+1) \cdots (z+n)}. \quad (18)$$

It is easy to prove (using the expression with the limit) that with this definition the recursion relation (2) is satisfied, and also that for $z = 1$ gives $\Gamma(1) = 1$.

Another one, due to Weierstrass, is

$$\Gamma(z) = \frac{e^{-\gamma z}}{z} \prod_{n=1}^{\infty} \frac{e^{z/n}}{1 + \frac{z}{n}} = e^{-\gamma z} \lim_{n \rightarrow \infty} \frac{n! \exp \left[z \left(1 + \frac{1}{2} + \dots + \frac{1}{n} \right) \right]}{z(z+1) \cdots (z+n)}, \quad (19)$$

where γ is the Euler-Mascheroni number defined in Section 2.2. The equivalence to the Euler definition can be seen by taking the logarithm of these two forms: the first gives

$$\log \Gamma(z) = -\log z + \sum_{n=0}^{\infty} \left[z \log \left(1 + \frac{1}{n} \right) - \log \left(1 + \frac{z}{n} \right) \right] \quad (20)$$

and the second gives

$$\log \Gamma(z) = -\log z - \gamma z + \sum_{n=0}^{\infty} \left[\frac{z}{n} - \log \left(1 + \frac{z}{n} \right) \right]. \quad (21)$$

Comparing these two expressions we note that they are equal because of eq. (48). Note that these two expressions are useful in order to numerically approximate $\log \Gamma(z)$, by truncating the series to a certain value of n .

1.3 Euler reflection formula

A useful formula is

$$\Gamma(z) \Gamma(1-z) = \frac{\pi}{\sin(\pi z)} \quad (22)$$

called the Euler reflection formula. To demonstrate it, consider the alternative definition (18) and note that

$$\begin{aligned} \frac{1}{\Gamma(z) \Gamma(1-z)} &= \frac{1}{-z \Gamma(z) \Gamma(-z)} \\ &= \frac{z(-z)}{-z} \prod_{n=1}^{\infty} \frac{(1 + \frac{z}{n})(1 - \frac{z}{n})}{(1 + \frac{1}{n})^z (1 + \frac{1}{n})^{-z}} \\ &= z \prod_{n=1}^{\infty} \left(1 - \frac{z^2}{n^2} \right) \\ &= \frac{\sin(\pi z)}{\pi} \end{aligned} \quad (23)$$

where we have used

$$\sin(\pi z) = \pi z \prod_{n=1}^{\infty} \left(1 - \frac{z^2}{n^2} \right). \quad (24)$$

1.4 Stirling approximation

The Stirling approximation for the factorial is

$$\boxed{n! = \sqrt{2\pi n} e^{-n} n^n} \quad (25)$$

which is valid for large n . To demonstrate it, we perform a saddle point approximation to the integral (1):

$$\begin{aligned} \Gamma(z+1) &= \int_0^\infty dt e^{-t} t^z \\ &= \int_0^\infty dt e^{-t+z \log t} \end{aligned}$$

now we expand the exponent near its maximum and get

$$\begin{aligned} &\simeq \int_0^\infty dt \exp\left(-z + z \log z - \frac{1}{2z} t^2\right) \\ &\simeq e^{-z} z^z \int_{-\infty}^\infty dt e^{-\frac{1}{2z} t^2} \\ &= \sqrt{2\pi z} e^{-z} z^z \end{aligned} \quad (26)$$

that is the claimed result.

1.5 Derivative recursion relation

Consider the Taylor expansion near $z = 0$ of $\Gamma(z + z_0 + 1)$:

$$\Gamma(z + z_0 + 1) = \sum_{k=0}^{\infty} \frac{\Gamma^{(k)}(z_0 + 1)}{k!} z^k ; \quad (27)$$

using eq. (2) this is equal to $(z + z_0) \Gamma(z + z_0)$, that expanded is

$$\begin{aligned} (z + z_0) \Gamma(z + z_0) &= (z + z_0) \sum_{k=0}^{\infty} \frac{\Gamma^{(k)}(z_0)}{k!} z^k \\ &= z_0 \Gamma(z_0) + \sum_{k=1}^{\infty} \frac{z^k}{k!} \left[z_0 \Gamma^{(k)}(z_0) + k \Gamma^{(k-1)}(z_0) \right] . \end{aligned} \quad (28)$$

Comparing the two expansions, we get the equality ($k > 0$)

$$\boxed{\Gamma^{(k)}(z_0 + 1) = z_0 \Gamma^{(k)}(z_0) + k \Gamma^{(k-1)}(z_0)} . \quad (29)$$

1.6 Inverse Gamma function

We can define an inverse Gamma function

$$\boxed{\Delta(z) \equiv \frac{1}{\Gamma(z)}} \quad (30)$$




Figure 2: $\text{Re } \Delta(x)$ for real x .

which is an entire function (analytical in all the complex plane). The recursion relation (2) becomes

$$\Delta(z) = z \Delta(z + 1) . \quad (31)$$

We are able to produce an integral representation of the Δ function: consider the Laplace transform of the function t^{z-1} with respect to t :

$$\int_0^{\infty} dt e^{-st} t^{z-1} = s^{-z} \Gamma(z) \quad (32)$$

and then we have that the inverse Laplace transform of s^{-z} is

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} ds e^{ts} s^{-z} = \Delta(z) t^{z-1} \quad (33)$$

where c has to be chosen to the right of all the singularities of s^{-z} , that is $c > 0$. Now, by choosing $t = 1$, we get

$$\Delta(z) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} ds e^s s^{-z} , \quad (34)$$

where the integration path can be deformed to be a contour of the negative real axis. Note that for positive integers $z = n + 1$ the branch cut vanishes and only a pole in $s = 0$ remains, and then the path can be closed around $s = 0$ to obtain

$$\frac{1}{n!} = \frac{1}{2\pi i} \oint ds e^s s^{-n-1} , \quad (35)$$

that can be also easily demonstrated using the residue theorem.

Now consider the Taylor expansion near $z = 0$ of $\Delta(z + z_0)$:

$$\Delta(z + z_0) = \sum_{k=0}^{\infty} \frac{\Delta^{(k)}(z_0)}{k!} z^k ; \quad (36)$$

using eq. (31) this is equal to $(z + z_0) \Delta(z + z_0 + 1)$, that expanded is

$$(z + z_0) \Delta(z + z_0 + 1) = (z + z_0) \sum_{k=0}^{\infty} \frac{\Delta^{(k)}(z_0 + 1)}{k!} z^k$$

$$= z_0 \Delta(z_0 + 1) + \sum_{k=1}^{\infty} \frac{z_0^k}{k!} \left[z_0 \Delta^{(k)}(z_0 + 1) + k \Delta^{(k-1)}(z_0 + 1) \right]. \quad (37)$$

Comparing the two expansions, we get the equality ($k \geq 1$)

$$\boxed{\Delta^{(k)}(z_0) = z_0 \Delta^{(k)}(z_0 + 1) + k \Delta^{(k-1)}(z_0 + 1)} \quad (38)$$

and, for $z_0 = 0$,

$$\Delta^{(k)}(0) = k \Delta^{(k-1)}(1). \quad (39)$$

1.6.1 Saddle point approximation

The analogous of the Stirling approximation can be done for the $\Delta(z)$ function. The result is:

$$\Delta(z) \simeq \sqrt{\frac{z}{2\pi}} e^z z^{-z}. \quad (40)$$

2 Gamma related functions

2.1 Log derivative $\psi(z)$

Define the logarithmic derivative of the Gamma function as follows

$$\boxed{\psi(z) \equiv \frac{d}{dz} \log \Gamma(z)} . \quad (41)$$

Form the recursion relation (2) it follows that

$$\boxed{\psi(z+1) = \psi(z) + \frac{1}{z}} , \quad (42)$$

as we can easily demonstrate:

$$\begin{aligned} \psi(z+1) &= \frac{d}{dz} \log \Gamma(z+1) \\ &= \frac{d}{dz} \log [z\Gamma(z)] \\ &= \frac{1}{z} + \psi(z) \end{aligned} \quad (43)$$

(note that the same result can be obtained from eq. (29) for $k = 1$). For a positive integer n , we can iterate the recursion (42) to obtain

$$\boxed{\psi(n+1) = \psi(1) + 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}} , \quad (44)$$

and the value of $\psi(1)$ is

$$\boxed{\psi(1) = -\gamma} \quad (45)$$

where γ is the Euler-Mascheroni number, as demonstrated in Section 2.2.

The functions

$$\boxed{\psi_n(z) \equiv \psi^{(n)}(z) = \frac{d^{n+1}}{dz^{n+1}} \log \Gamma(z)} \quad (46)$$

are also defined. Iterating the derivative in (42) we obtain the recursion relation

$$\boxed{\psi_n(z+1) = \psi_n(z) + (-1)^n n! \frac{1}{z^{n+1}}} . \quad (47)$$

2.2 Euler-Mascheroni number

The Euler-Mascheroni number is defined as

$$\boxed{\gamma \equiv \sum_{n=1}^{\infty} \left[\frac{1}{n} - \log \left(1 + \frac{1}{n} \right) \right]} , \quad (48)$$

and its value is

$$\boxed{\gamma = 0.577216\dots} . \quad (49)$$

Note that

$$\begin{aligned}
\sum_{n=1}^{\infty} \log\left(1 + \frac{1}{n}\right) &= \lim_{N \rightarrow \infty} \sum_{n=1}^N \log\left(1 + \frac{1}{n}\right) \\
&= \lim_{N \rightarrow \infty} \sum_{n=1}^N [\log(1+n) - \log n] \\
&= \lim_{N \rightarrow \infty} \log(1+N) \\
&= \lim_{N \rightarrow \infty} \log N, \tag{50}
\end{aligned}$$

where we noted that the sum is a telescopic sum and that in the limit $N \rightarrow \infty$ $\log(1+N)$ and $\log N$ are equivalent. Thus we can rewrite eq. (48) as

$$\boxed{\gamma = \lim_{N \rightarrow \infty} \left[\sum_{n=1}^N \frac{1}{n} - \log N \right]}. \tag{51}$$

Now we prove that $\boxed{\psi(1) = -\gamma}$. To do this, let us consider eq. (20), and take its derivative:

$$\psi(z) = -\frac{1}{z} + \sum_{n=1}^{\infty} \left[\log\left(1 + \frac{1}{n}\right) - \frac{1}{n} \frac{1}{1 + \frac{z}{n}} \right]. \tag{52}$$

For $z = 1$ we get

$$\begin{aligned}
\psi(1) &= -1 + \sum_{n=1}^{\infty} \left[\log\left(1 + \frac{1}{n}\right) - \frac{1}{1+n} \right] \\
&= -\sum_{n=1}^{\infty} \left[\frac{1}{n} - \log\left(1 + \frac{1}{n}\right) \right], \tag{53}
\end{aligned}$$

in which we recognise the definition (48).

2.3 Generalized Gamma functions

You can also define an incomplete Gamma function (or plica function)

$$\Gamma(z, a) \equiv \int_a^{\infty} dt e^{-t} t^{z-1} \tag{54}$$

and a truncated Gamma function

$$\gamma(z, a) \equiv \int_0^a dt e^{-t} t^{z-1}, \tag{55}$$

that have a branch cut on the negative real axis in the a complex plane. Obviously

$$\Gamma(z) = \Gamma(z, a) + \gamma(z, a). \tag{56}$$

For integer $z = k + 1$, integrating repeatedly by parts we get

$$\Gamma(k + 1, a) = k! e^{-a} \sum_{n=0}^k \frac{a^n}{n!} \quad (57)$$

$$\gamma(k + 1, a) = k! \left(1 - e^{-a} \sum_{n=0}^k \frac{a^n}{n!} \right). \quad (58)$$

2.4 Beta function

The Beta function is defined as

$$B(a, b) \equiv \int_0^1 dx x^{a-1} (1-x)^{b-1} = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}. \quad (59)$$

To show the last equality, consider the product

$$\begin{aligned} \Gamma(a)\Gamma(b) &= \int_0^\infty dt e^{-t} t^{a-1} \int_0^\infty du e^{-u} u^{b-1} \\ &= 4 \int_0^\infty dx e^{-x^2} x^{2a-1} \int_0^\infty dy e^{-y^2} y^{2b-1} \\ &= \int_{-\infty}^\infty dx \int_{-\infty}^\infty dy e^{-(x^2+y^2)} |x|^{2a-1} |y|^{2b-1} \\ &= \int_0^{2\pi} d\theta |\cos^{2a-1} \theta \sin^{2b-1} \theta| \int_0^\infty dr e^{-r^2} r^{2a+2b-1} \\ &= 2 \Gamma(a+b) \int_0^{\pi/2} d\theta \cos^{2a-1} \theta \sin^{2b-1} \theta \\ &= \Gamma(a+b) \int_0^1 dx x^{a-1} (1-x)^{b-1}. \end{aligned} \quad (60)$$