

Paths to Geant4 evolution: Refactoring, Reengineering and Physics

State-of-the-Art Simulation of Photon Interactions with Matter

Maria Grazia Pia



M. Batič, M. Han, S. Hauf, G. Hoff, C. H. Kim, M. Kuster,
P. Saracco, H. Seo, G. Weidenspointner, A. Zoglauer



INFN Genova, Italy
J. Stefan Institute, Ljubljana, Slovenia
Hanyang University, Seoul, Korea
PUCRS, Porto Alegre, Brazil
EU XFEL GmbH, Hamburg, Germany
MPI Halbleiterlabor, Munich, Germany
Space Sciences Laboratory, UC Berkeley, USA

NSS 2012
30 October 2012, Disneyland, USA





"The code slowly sinks from engineering to hacking".

“Refactoring is the process of changing a software system in such a way that it does not alter the external behavior of the code yet improves its internal structure.”

“When you refactor you are improving the design of the code after it has been written.”


Reengineering “seeks to transform a legacy system into the system you would have built if you had the luxury of hindsight and could have known all the new requirements that you know today.”

“A legacy is something valuable that you have inherited.”

“With rapid development tools *and* rapid turnover in personnel, software systems can turn into legacies more quickly than you might imagine.”

R&D Project

Series of pilot projects
going on since 2008



Focus on **physics** and
fundamental concepts
of particle transport

Highlights
No time to go into details


State-of-the-art, quantified

Geant 4



Archival
literature

Produce concrete deliverables





Smells


If it stinks, change it.

Grandma Beck, discussing child-rearing philosophy

M. Fowler, K. Beck et al.,

Refactoring: Improving the Design of Existing Code

- Duplicated Code
- Long Method
- Large Class
- Long Parameter List
- Divergent Change
- Shotgun Surgery
- Feature Envy
- Data Clumps
- Primitive Obsession
- Switch Statements
- Parallel Inheritance Hierarchies
- Lazy Class
- Speculative Generality
- Temporary Field
- Message Chains
- Middle Man
- Inappropriate Intimacy
- Alternative Classes with Different Interfaces
- Incomplete Library Class
- Data Class
- Refused Bequest



Electromagnetic smells

Coupling

σ_{tot} and final state modeling have been decoupled in hadronic physics design since RD44

Dependencies

on other parts of the software

“model”

Total cross section

Whether a process occurs

Final state generation

How a process occurs




One needs a geometry
(and a full scale application)
to test (verify) a cross section

Difficult to test → no testing
often

Problem domain analysis
Improve domain decomposition






Benefits

Transparency
Ease of maintenance
Simplicity of testing for V&V

Basic physics V&V can be performed by means of **lightweight unit tests**

Easier exploration of new models
Easier quantification of accuracy

Bare-bone physical functionality



Decorations on top

Motion of atomic electrons
Doppler broadening
Binding effects

scattering functions

Vacancy creation
Atomic relaxation

Doppler profiles



Prune

Number one in the stink parade is duplicated code physics

M. Fowler,
Refactoring

Objective quantification of smell

Two Geant4 models, identical underlying physics content (*it used to be different*)



Efficiency w.r.t. experiment

“Livermore”	Penelope
EPDL97	EPDL97
0.38 ± 0.06	0.38 ± 0.06

Code bloat

Burden on

- Software design
- Maintenance
- User support

Unnecessary complexity

Trash and redo

Number one in the stink
parade is duplicated code

1. Bearden & Burr (1967)
2. Carlson
3. EADL
4. Sevier
5. Tol 1978 (Shirley)
6. Tol 1996 (Larkins)
7. Williams


numbers

Atomic binding energies

Geant4 { Carlson + Williams
EADL



(Carlson
Shirley)

Source of epistemic uncertainties?



23

pages



Algorithms

Popular belief

Physics model X is intrinsically slow

Baroque methods to combine it with “faster” lower precision models and limit its use to cases where one is willing to pay for higher precision


This design introduces an additional computational burden due to the effects of inheritance and the combination algorithms themselves

Truth

Physics model X is intrinsically fast

But its computationally fast physics functionality is spoiled by an inefficient sampling algorithm

Change the sampling algorithm!



- ▶ No code smell
- ▶ Spotted through in-depth **code review** in the course of **software validation**

The fastest algorithm

no algorithm at all

Shift modeling from algorithms to data

Merging models




Smoothing data

Guidance from
experimental data
(*when available*)


Data libraries

- PIXE ($\sigma_{\text{ionisation}}$: p)
- BEB–DM ($\sigma_{\text{ionisation}}$: e^-)
- Photon elastic
 - RTAB (L. Kissel)

Electron impact ionisation cross sections



...no silver bullet



Photon elastic scattering cross section at 90°

*Experimental data by Schumacher et al.,
Starr et al., Jackson et al., Moreh et al. etc.,
complete list in related publication*

Smoothing models (or data) that exhibit significantly different behaviour is not physically justified

← Here our understanding of the underlying physics phenomena fails

Experimental data themselves must be analyzed for inconsistencies and possible systematics

How to solve this kind of problems?


Better theoretical calculations
New, dedicated experimental data

No easy solution

Refactoring data management


- Today's technology
 - ...keeping an eye on the new C++ Standard

- Optimal container
- Pruning data
- Splitting files
- Software design




Big refactoring

Well defined
responsibilities
and interactions




Geant4 Radioactive Decay






Experiment: Z. W. Bell (ORNL)

Validation




Statistical Decay



More in
N28-6/7

New algorithm

Performance



Paths to Geant4 evolution: Refactoring, Reengineering and Physics

State-of-the-Art Simulation of Photon Interactions with Matter

Maria Grazia Pia

M. Batič, M. Han, S. Hauf, G. Hoff, C. H. Kim, M. Kuster,
P. Saracco, H. Seo, G. Weidenspointner, A. Zoglauer



INFN Genova, Italy
J. Stefan Institute, Ljubljana, Slovenia
Hanyang University, Seoul, Korea
PUCRS, Porto Alegre, Brazil
EU XFEL GmbH, Hamburg, Germany
MPI Halbleiterlabor, Munich, Germany
Space Sciences Laboratory, UC Berkeley, USA

NSS 2012
30 October 2012, Disneyland, USA



Refactoring to improve physics

Refactoring

M. Fowler, Refactoring

is the process of changing a software system in such a way that
it **does not alter the external behavior** of the code
yet improves its internal structure

Refactoring/reengineering legacy simulation code is an opportunity

- To complement software testing with **physics validation**
- To evaluate and compare the **accuracy** of various modeling options
- To estimate the trade-off between **accuracy** and **performance**
- To document the **state-of-the-art** for Monte Carlo transport

Electron-photon physics

Models and data libraries used by Geant4 and other
Monte Carlo codes



Photon elastic scattering

Physics

- Rayleigh scattering
- Thomson nuclear scattering
- Delbrück scattering
- Nuclear resonance scattering

Form factor approximation

- **Non relativistic** form factors
(Hubbell 1977) **FLUKA, Geant4, MCNP, Penelope**
- **Relativistic** form factors
(Hubbell & Øverbø) **EGS, ITS, XCOM**
- **Modified** form factors
(Schaupp *et al.*)
- + **anomalous scattering** factors

Geant 4

“Livermore”	EPDL
“Penelope 2008”	EPDL
“Penelope 2001”	Baro <i>et al.</i>
“Standard”	Unknown


*EPDL: non relativistic form factors
Thomson scattering in total cross
section only*

2nd order S-matrix

- Calculations by L. Kissel
- Not yet exploited in Monte Carlo codes

Forward peaked

Heavy materials




Comparison with experiment

> 4000 differential cross section measurements from the literature


$$5.41 \text{ keV} \leq E \leq 39 \text{ MeV}$$

$$0.5^\circ \leq \theta \leq 165^\circ$$

$$69 \text{ target atoms}, 3 \leq Z \leq 92$$







MFASF
RFASF
MF
RF
NF
Penelope2001
Penelope2008
EPDL
SM



Statistical analysis of compatibility + categorical analysis

Nuclear Thomson scattering

Important to account for it especially at higher energy



Model validation

State of the art

Identified through quantitative validation w.r.t. experiment

Quantification

Compatibility with experiment: **GoF test**
(χ^2 test)

Comparison of models' accuracy: **categorical analysis**
(Fisher's exact test and χ^2 test)

Differential cross sections

	Penelope 2001	Penelope 2008	EPDL	Relativ. FF	Non-Rel. FF	Modified FF	MFF ASF	RFF ASF	SM NT
ϵ	0.27	0.38	0.38	0.25	0.35	0.49	0.52	0.48	0.77
error	± 0.05	± 0.06	± 0.06	± 0.05	± 0.06	± 0.06	± 0.06	± 0.06	± 0.05

ϵ = fraction of test cases compatible with experiment
0.01 significance in the χ^2 test




S-matrix

Modified form factors




Other physics
to be
accounted for
at higher
energy



Total elastic cross section

- S-matrix
- EPDL
- MFASF
- XCOM
- Storm & Israel
- Brennan & Cowan
- Geant4 Standard

Scarce experimental data





Photoelectric effect

Total cross section

- EPDL
- Kissel (RTAB)
- Storm & Israel
- Biggs & Lighthill
- XCOM
- Brennan & Cowan
- McMaster

Total cross section
Shell cross sections
Photoelectron angular distribution

Geant 4

“Livermore”

“Penelope 2008”

“Penelope 2001”

“Standard”

EPDL

EPDL



EPDL

Biggs &
Lighthill

XCOM?



EPDL used down to 250 eV
Never validated, nor optimized

Total cross section difference w.r.t. experiment



Geant4

“Livermore” and “Penelope”:
both EPDL cross sections



Inadequate modeling or systematics in the measurements?

Photoelectric effect

Total cross section

$E > 50 \text{ keV}$

$E=2.62 \text{ MeV}$




$E = 661.6 \text{ keV}$

$E = 279.2 \text{ keV}$

$E = 123.6 \text{ keV}$

$E = 59.54 \text{ keV}$

PRELIMINARY




Photoelectric effect

Total cross section


100 eV - 1 keV

Small experimental sample
Noble gases


$Z = 2$




$Z = 10$




$Z = 18$



$Z = 36$



$Z = 54$



500 eV – 1 keV

p-value_{EPDL} = 0.593

250 – 500 eV

p-value_{EPDL} = 0.029

PRELIMINARY


Photoelectric effect

PRELIMINARY

Total cross section

10-100 eV

Measured
in gases
only



Systematics in
experiment?

Inadequate
calculations?

Conclusions

A quantitatively validated, state-of-the-art
Monte Carlo code



Large effort

Supported by software design


a tremendous challenge
an inspiring motivation

Refactoring techniques and reengineering patterns contribute
to **computational performance** and facilitate **validation**

Quantitative validation of physics models

more about it in the last talk of this session

Physics insight is the key



...a big **THANK YOU**
to the CERN Library!