



Physics methods for the simulation of photoionisation

T. Basaglia¹, M. Batic², M. C. Han³, G. Hoff⁴,
C. H. Kim³, H. S. Kim³, M. G. Pia⁵, P. Saracco⁵

¹CERN

²Sinergise, Ljubljana, Slovenia

³Hanyang University, Seoul, Korea

⁴Pontificia Universidade Catolica do Rio Grande do Sul, Porto Alegre, Brazil

⁵INFN Genova, Italy

IEEE NSS 2013

27 October – 2 November 2013

Seoul, Korea

2013 IEEE NSS/MIC/RTSD
"Beyond Imagination of Future Science"

Rationale

Simulation physics

The simulation of photon physics is well-established.

→ Is there any quantitative validation?

New physics

New theoretical calculations and parameterisations were released recently.

→ Are they accurate than old one?

New trends

Very low energy(~eV scale) & micro/nano dosimetry
ex. FLUKA, Geant4-DNA, MOCA, OREC/NOREC, PARTRAC, Penelope, PTB-code, Trion etc.

→ How accurate are models?

Project to validate **quantitatively** a wide set of simulation modeling options against a large collection of experimental data



State-of-the-art simulation of photon interactions

Elastic scattering
: Published


Photoelectric effect

Compton scattering
pair production
: first results

Photoionisation in Monte Carlo codes

| | Cross sections | Angular distribution |
|-----------------|--|----------------------------------|
| EGS5 | PHOTX | Sauter |
| EGSnrc | Storm-Israel Fit to XCOM EPDL97 (subshell) | Sauter |
| FLUKA | EPDL97 | Sauter |
| Geant4 | Revised Biggs-Lighthill (Henke) EPDL97 | Sauter-Gavrila Same direction |
| ITS | Scofield 1973 | Fischer+Sauter |
| MCNP(X) | EPDL89, EDPL97 ENDFB/IV+Storm-Israel | |
| Penelope | EPDL97 | Sauter (K shell) |

Photoionisation in Geant4 9.6



Cross section sources

| Year | Compilation | Energy | Z | (sub)Shell | Method |
|-----------|--------------------------|------------------|-------|------------|---------------|
| 1967-1988 | Biggs-Lighthill | 10 eV – 100 GeV | 1-100 | - | parameterised |
| 1992 | Brennan-Cowan | 30 eV – 700 keV | 3-92 | - | tabulated |
| 2000 | Chantler | 10 eV – 433 keV | 1-92 | K | tabulated |
| 2003 | Ebel | 1 keV – 300 keV | 1-92 | all | parameterised |
| 2002 | Elam | 100 eV – 1 MeV | 1-98 | - | tabulated |
| 1997 | EPDL97 (Scofield) | 10 eV – 100 GeV | 1-100 | all | tabulated |
| 1982-1993 | Henke | 10 eV – 30 keV | 1-92 | - | tabulated |
| 1970-2006 | McMaster/Shaltout | 1 keV – 700 keV | 1-94 | - | tabulated |
| 1989 | PHOTX (Scofield) | 1 keV – 100 MeV | 1-100 | | tabulated |
| 2001 | RTAB | 10 eV – 30 keV | 1-99 | all | tabulated |
| 1973 | Scofield | 1 keV – 1.5 MeV | 1-100 | all | tabulated |
| 1970 | Storm-Israel | 1 keV – 100 GeV | 1-100 | - | tabulated |
| 1973 | Veigle | 100 eV – 100 MeV | 1-94 | - | tabulated |
| 1987-2010 | XCOM (Scofield) | 1 keV – 100 GeV | 1-100 | - | tabulated |

Different methods and calculations

e.g. Chantler's exchange potential in his DHF calculation is different from Scofield's


Strategy

- Evaluate a large number of available modeling options
- Suitable for use in Monte Carlo simulation codes
 - Tabulated theoretical calculations
 - Simple analytical formulations, with documented parameters
- All options evaluated in the same computational environment
 - Minimize dependencies on other software parts (*not always components*)
- Quantitative, objective evaluation based on **statistical methods**
- Establish state-of-the-art for the simulation of photoionisation on objective ground
- Computational performance measured along with physical accuracy

※ We only focus on the simulation with non-polarised photons case.

Computational environment


Streamlined software design consistent with Geant4 kernel



Sharp domain decomposition
Clearly identified responsibilities
No duplication of code nor of functionality

Policy-based class design
(à la Alexandrescu, *Modern C++ design*, 2001)

- minimize dependencies
- lightweight unit tests for validation



Experimental data

Collected from the literature

- Total cross sections
- Partial cross sections
- Angular distributions

> 150 references
> 5000 data points
~ 3700 σ_{total}
~ 1400 σ_{shell}



Data types

- **Pure experimental** cross sections: direct measurements
- **Semi-empirical** cross sections: involve theoretical manipulations
 - ▷ e.g. subtraction of calculated scattering contribution (Compton and elastic)


Format

- Tables, text
- Figures: digitized, digitization error estimated

Evaluation of experimental data

- Systematic effects: identified whenever possible
- Outliers

Systematic effect?



Difference between calculated and “experimental” total cross sections, expressed in terms of number of standard deviations:
pure experimental
 and
semi-empirical
 data

Only pure experimental data used in the validation process

Data analysis method

- Two-stage statistical analysis
 1. Compatibility of each cross section calculation method with experiment
 2. Comparison of compatibility with experiment across modeling categories
- Quantitative appraisal of capabilities and differences

Compatibility with experiment

Goodness-of-fit test

χ^2 test

$\alpha = 0.01$

$\alpha \geq 0.01 \rightarrow$ pass

$\alpha < 0.01 \rightarrow$ fail

Difference across categories

Contingency tables

Fisher exact test

Barnard test


Pearson χ^2 test

$\alpha = 0.05$

efficiency= $N_{\text{pass}} / N_{\text{test cases}}$

as appropriate

Results - Total cross sections




- Most calculation methods exhibit similar compatibility with experiment for **E>250 eV**
 - Chantler, Brennan-Cowan look worse*
- Degraded accuracy below 250 eV


Analysis of contingency tables

| | EPDL Chantler | EPDL Brennan-Cowan |
|--|------------------|-----------------------|
|--|------------------|-----------------------|

| | |
|------------------|-------|
| Fisher | 0.044 |
| Pearson χ^2 | 0.033 |
| Barnard | 0.035 |



Results - Shell cross sections



Systematic effect observed with
RTAB shell cross sections
(presumably a missing factor in the calculation)

Calculated **inner shell** cross sections compatible with experiment

Outer shell cross sections inconsistent with experimental data
Beware: small data sample, limited data sources





p-value χ^2 test

| shell | EPDL | Chantler | RTAB | scRTAB | Ebel |
|-------|--------|----------|--------|--------|--------|
| K | 0.209 | 0.350 | <0.001 | 0.315 | <0.001 |
| L1 | 0.075 | | <0.001 | 0.069 | 0.964 |
| L2 | 0.339 | | <0.001 | 0.299 | 0.154 |
| L3 | 1 | | <0.001 | 1 | 1 |
| M1 | <0.001 | | <0.001 | <0.001 | |
| M4 | 0.031 | | <0.001 | <0.001 | |
| M5 | <0.001 | | <0.001 | <0.001 | |
| N1 | <0.001 | | <0.001 | <0.001 | |
| N6 | <0.001 | | <0.001 | <0.001 | <0.001 |
| N7 | <0.001 | | <0.001 | <0.001 | <0.001 |
| O1 | <0.001 | | <0.001 | <0.001 | <0.001 |
| O2 | <0.001 | | <0.001 | <0.001 | <0.001 |
| O3 | <0.001 | | <0.001 | <0.001 | <0.001 |
| P1 | <0.001 | | <0.001 | <0.001 | <0.001 |

Results - Angular distribution

Option à la GEANT 3 (Sauter) evaluated along with other Geant4 options


Qualitative appraisal
Limited experimental sample
Experimental systematic effects
(corrected/uncorrected data)



Conclusion

- **Large scale effort to evaluate quantitatively physics methods for photoionisation simulation**
 - Part of a wider project for **quantitative** assessment of **state-of-the-art** simulation of photon interactions
- **Total cross section**
 - Most calculation methods exhibit similar behaviour
 - More recent calculations (Chantler, Brennan-Cowan) do not appear more accurate than old Scofield's 1973 (unrenormalized)
- **Inner shells**
 - EPDL, (*corrected*) RTAB appear equivalent, Ebel's parameterisation inconsistent with experimental K shell data
- **Outer shells**
 - No calculation method appears adequate to reproduce experimental data
- **Photoelectron angular distribution**
 - Scarce data and experimental systematics prevent a quantitative discrimination

All results will be documented in detail in a forthcoming publication



...a big **THANK YOU** to
the CERN Library!