

$\gamma p \rightarrow K_s^0 \Sigma^+$ photoproduction

- ◆ Motivation
- ◆ Experiment
- ◆ Results
- ◆ Interpretation
- ◆ Summary & Outlook

Crystal-Barrel/TAPS
BGO-OD collaborations

Hartmut Schmieden

Physikalisches Institut
Universität Bonn

SFB/TR-16




Motivation

- ❖ relevant degrees of freedom ?
- ❖ dynamics behind baryon excitation spectrum ?
 - 3q
 - 3q-g hybrids
 - q-2q quark-diquark correlations
 - qqq-q \bar{q} baryon-meson
- ❖ $|N\rangle = |\text{qqqq}\rangle + |\text{qqqq q}\bar{q}\rangle + |\text{qqqq g}\rangle + \dots$
 - $|\text{qqqq}\rangle$ short range
 - $|\text{qqqq q}\bar{q}\rangle$ long range
 - $|\text{qqqq g}\rangle$ low t

Motivation

Why $K^0\Sigma^+$ channel ??




- ◆ non-pionic **photoproduction** of baryon resonances
- ◆ Born contributions suppressed compared to K^+ production
- ◆ in particular: t-channel
- ◆ but: K^{0*} exchange
- ◆ box-formation possible



Motivation


Why $K^0\Sigma^+$ channel ??

- ◆ non-pionic **photoproduction** of baryons
- ◆ Born contributions suppressed compared to pion exchange
- ◆ in particular: t-channel
- ◆ but: K^{0*} exchange
- ◆ box-formation possible



$\gamma p \rightarrow K^0\Sigma^+ @ K^* \text{ threshold}$

previous data inconclusive






ELSA experimental areas


Beamline E

Beamline S




© D. Elsner

Experiment: Crystal Barrel / TAPS




Event reconstruction & data analysis




kinematic fit

- beam E_γ along axis
- target proton at rest
- detected γ 's E, θ, ϕ
- ejected p not required




Event reconstruction & data analysis

acceptance



$\gamma + p \rightarrow K^0 + \Sigma^+$: cross sections

R. Ewald et al., PLB 713 (2012)



this work (CB polarised beam)


Castelijns (CB unpol. beam, EPJ A35 (08) 39)
SAPHIR, R. Lawall et al., EPJ A24 (05) 275


main syst. errors

- backgrd subtr.
- kin.fit (cut cf level)
- acceptance
- photon flux

$\gamma + p \rightarrow K^0 + \Sigma^+$: cross sections


R. Ewald et al., PLB 713 (2012)






total cross section

R. Ewald et al., PLB 713 (2012)



total cross section


R. Ewald et al., PLB 713 (2012)




Interpretation

BoGa PWA

A.V. Anisovich et al., Phys. Lett. B 711 (2012) 162




- resonances indicated below 2000 MeV
 $N_{1/2^-}$ (1895) & $N_{3/2^-}$ (1875)
- → parity doublets with
 $N_{1/2^+}$ (1875) & $N_{3/2^+}$ (1900)
- E. Klemp's idea:
“missing” res's
↔ complete SU(6) multiplets ??



Interpretation


A. Ramos & E. Oset, arXiv: 1304.7975

- vectormeson-baryon dynamics
- structure \leftrightarrow $N^*(2080)(3/2^-)$ / $N^*(2090)(1/2^-)$ [removed from PDG]
- delicate interference $K^*\Lambda$ / $K^*\Sigma$ channels
- $\rightarrow K^0\Sigma^0$ off neutron target
- ω photoproduction ??
- K^0 beam asymmetry ??



Results: Photon Beam Asymmetry


Nam, Choi, Hosaka, Kim, Phys. Rev. D75 (2007) 014027




$$\Sigma = \frac{\sigma_{\perp} - \sigma_{\parallel}}{\sigma_{\perp} + \sigma_{\parallel}}$$

Results: Photon Beam Asymmetry

Nam, Choi, Hosaka, Kim, Phys. Rev. D75 (2007) 014027




Results: Photon Beam Asymmetry




ω photoproduction: beam-target asymmetry

H. Eberhardt, Dr. thesis (2012)




energy of x-sec anomaly
in $K^0\Sigma^+$ channel



BGO-OD experiment @ ELSA

- low-t to investigate meson-baryon components
- → forward angles / acceptance
- BGO-OD setup
 - high-res. forward detector
 - hermetic crystal calorimeter
- complementary to CB, CLAS, LEPS

$$\begin{aligned} K^0_S &\rightarrow \left\{ \begin{array}{ll} \pi^0 \pi^0 & 31 \% \\ \pi^+ \pi^- & 67 \% \end{array} \right. \\ \omega &\rightarrow \left\{ \begin{array}{ll} \pi^0 \gamma & 8.5 \% \\ \pi^+ \pi^- \pi^0 & 89 \% \end{array} \right. \end{aligned}$$



BGO-OD experiment

TOF walls

Horizontal & vertical bars 20cm wide, 3m long
Time res < 1ns

Drift Chambers

8 chambers, 2 layers each
x, y, u (+9 deg), v (-9 deg)
8.5 mm drift cell radius

SciFi


2 layers of 288 (x) and 352 (y) fibres,
3mm diameter, 90 deg rotation
66 x 51 cm, $\Delta t \sim 2\text{ns}$

MOMO

3 layers of 244 fibres,
60 deg rotation.
Spatial resolution $\sim 1.5\text{mm}$

BGO Ball

480 $\text{Bi}_4\text{Ge}_3\text{O}_{16}$ crystals
25 – 155 degrees
Internal plastic scintillator
& MWPC
for particle tracking



Forward spectrometer

+/- 8 deg (vertical), +/ - 10 deg (horizontal) acceptance

IH₂ target

Centre of BGO Ball





Photon Tagger

120 channels, 10-90% of e- energy spectrum - linear and circularly polarised photons

ELSA e- beam

< 3.5 GeV

reconstruction: BGO ball



Conclusions

- ◆ photoproduction of $K^0_S \Sigma^+$ off protons
- ◆ differential and total cross sections
with full angular coverage & flat acceptance [R. Ewald et al., PLB 713 (2012)]
- ◆ beam asymmetry Σ (and recoil polarisation)
- ◆ indication for “cusp-like“ structure in cross section
in forward directions between $K^* \Lambda / \Sigma$ thresholds
- ◆ visible also in total cross section
- ◆ smooth transition t- / s-channel
- ◆ dominant $1/2^-$ & $3/2^-$ waves in PWA
- ◆ **K* - Y quasibound state ?** \leftrightarrow hint also from beam asymmetry
- ◆ dynamically generated states: vectormeson nonet \times baryon octet
 - Oset & Ramos, EPJ A 44 (2010) 445
 - Ramos & Oset, arXiv:1304.7975
 - ↳ K^0 off neutron target !
 - ↳ photon beam asymmetry
 - ↳ ω channel

THANKS

DFG SFB/TR-16

Federal State of Northrhine-Westphalia

data analysis: doctoral thesis [Ralf Ewald](#)

Oliver Jahn & Tom Jude