

# A Forward Photon Tagging Facility for CLAS12

**M.Battaglieri**

*Istituto Nazionale di Fisica Nucleare  
Genova - Italy*

## CLAS12 EUROPEAN WORKSHOP

25-28 February 2009  
Genova, Italia

First European Workshop  
on the physics program and  
equipment of the CLAS12  
experiment at Jefferson Lab


**Location:**  
Dipartimento di Fisica dell'Università di Genova  
Via Dodecaneso 33, 16146 Genova, Italia

**Information:**  
<http://www.ge.infn.it/~clas12>  
clas12@ge.infn.it






# From CEBAF at 6 GeV






# From CEBAF at 6 GeV to CEBAF at 12 GeV



# The Hall-B photon tagger



- Gold and diamond radiator for In/Coherent Bremsstrahlung
- Energy coverage:  $0.2\text{-}0.95 E_0$
- Efficiency  $\sim 80\%$
- Energy Resolution  $\sim 10^{-3}$
- Timing Resolution  $\sim 100 \text{ ps}$



The existing dipole magnet is unable to deflect the 11 GeV primary beam on the existing beam-dump

The existing PHOTON TAGGER will be available for energies up to  $E_\gamma \sim 6.1 \text{ GeV}$

Options for  $E_\gamma > 6 \text{ GeV}$ ?

# Why photoproduction?

## Physics motivations

### Meson spectroscopy

Standard PWA on H target

Spectroscopy on He4 and other gas targets

### Hadron spectroscopy

Heavy mass baryon resonances (Cascades)

double-strangeness sets a higher mass

small width helps to detect and study excited states

### Compton scattering

### Meson polarizabilities

### J/ $\Psi$ production close threshold and on nuclear targets

### Large $-t$ physics

.....

# Why photoproduction?

## Physics motivations

### Meson spectroscopy

Standard PWA on H target

Spectroscopy on He4 and other gas targets

### Hadron spectroscopy

Heavy mass baryon resonances (Cascades)

double-strangeness sets a higher mass

small width helps to detect and study excited states

### Compton scattering

### Meson polarizabilities

### J/ $\Psi$ production close threshold and on nuclear targets



### Large $-t$ physics

.....

# Meson spectroscopy with photons at JLab


## Why photoproduction?

- ★ Photoproduction: exotic  $J^P$  are more likely produced by  $S=1$  probe



- ★ Production rate for exotics is expected comparable as for regular mesons



Few data (so far) but expected similar production rate as regular mesons



# Partial Wave Analysis

## 1) the isobar model

e.g.  $3\pi$  system



**Does the PWA work with photo-production data?**

**Use the PWA machinery on CLAS data**

## 2) Moments+Dispersion relations e.g. $2\pi$ system


### 1) Moments of the angular distribution in term of partial waves

$$\langle Y_{\lambda\mu} \rangle(E_\gamma, t, M) = \frac{1}{\sqrt{4\pi}} \int d\Omega_\pi \frac{d\sigma}{dt dm d\Omega_\pi} Y_{\lambda\mu}(\Omega_\pi)$$

$$\begin{aligned} \langle Y_{00} \rangle = N & [ |S|^2 + |P_-|^2 + |P_0|^2 + |P_+|^2 + |D_-|^2 \\ & + |D_0|^2 + |D_+|^2 + |F_-|^2 + |F_0|^2 + |F_+|^2 ] \end{aligned}$$

2) Parametrize partial waves in term of known  $\pi\pi$  phase shift and unknown coefficients using Dispersion Relations

3) Derive partial wave cross sections to compare with models




# Partial Wave Analysis with CLAS Isobar Model

$$\gamma p \rightarrow (n) \pi^+ \pi^+ \pi^-$$


★ Possible evidence of exotic meson  $\pi_1(1600)$  in  $\pi^- p \rightarrow p \pi^- \pi^- \pi^+$  (E852-Brookhaven)

★ Not confirmed in a re-analysis of a higher statistic sample

Simple final state with low background



## CLAS/g6c (preliminary)



★ Clear evidence of non-exotic  $2^{++}$  state  $a_2(1320)$

No-evidence of exotic  $1^+$  state  $\pi_1(1600)$

★ Relevance of baryon resonance background

PWA in CLAS is feasible!

# Partial Wave Analysis with CLAS

## Moments + Dispersion relations

$\gamma p \rightarrow p \pi^+ \pi^-$

$M(\pi^+\pi^-)$  spectrum below 1.5 GeV

P-wave:  $\rho$  meson






D-wave:  $f_2(1270)$

S-wave:  $\sigma$ ,  $f_0(980)$  and  $f_0(1320)$

### Fitted moments

$3.4 \text{ GeV} < E_\gamma < 3.6 \text{ GeV}$

$0.5 \text{ GeV}^2 < -t < 0.6 \text{ GeV}^2$



★ Known states are well reproduced, e.g.  $\rho(770)$

★ First observation of direct production of  $f_0(980)$  ( $\pi\pi$  S-wave) meson

Accepted by Phys.Rev.Lett. 2009

# Photoproduction experiments at JLab-12


## ★ The Detectors

- Determination of  $J^P$  of meson states requires **Partial Wave Analysis**
- Decay and Production of **exclusive** reactions
- Good acceptance, energy resolution, particle Id




## Hermetic charged/neutral particles detector

### Hall-D - GlueX Detector




### Hall-B - CLAS12 Detector



# Photoproduction experiments at JLab-12GeV

## \* The photon beam

- With a 11-12 GeV electron beam only few choices:
  - 1) Bremsstrahlung
  - 2) Quasi-real electro-production
- Tagger (initial photon energy) is required to add 'production' information to decay
- Linear polarization is useful to simplify the PWA and essential to isolate the nature of the t-channel exchange




- ★ Essential to isolate production mechanisms (M)
- ★ Polarization acts as a  $J^{PC}$  filter if M is known
- ★ Linear polarization separates natural and unnatural parity exchange

Hall-D and Hall-B will host real photon beam!


# Meson spectroscopy with photons at JLab-12GeV

## Coherent tagged Bremsstrahlung Hall-D



### Performance

- ★ (.5-.95)  $E_{beam} \rightarrow 6 < E_\gamma < 11$  GeV (10MeV resolution)
- ★ Photon Flux  $\sim 10^7 - 10^8$   $\gamma/s$
- ★ 30cm LH target  $\rightarrow L \sim 10^{31}$   $cm^{-2}s^{-1}$
- ★ Linear polarization  $\sim 50\% - 15\%$  (collective)



## Quasi-real electroproduction at very Low $Q^2$ Hall-B

$E_{scattered}$	1 - 4 GeV
$\theta$	$0.5^\circ - 1.2^\circ$
$\phi$	$0^\circ - 360^\circ$
$\nu$	7 - 10 GeV
$Q^2$	0.003 - 0.029 $GeV^2$
$W$	3.9 - 4.6 GeV
$x_{Bj}$	0.0001 - 0.002


### Performance

- ★  $7 < E_\gamma < 10$  GeV
- ★ 5cm LH target  $\rightarrow L \sim 10^{34}$   $cm^{-2}s^{-1}$
- ★ Linear polarization  $\sim 65\% - 20\%$  (individual)
- ★ Capability of forward tagging (electron detection)

# Real and quasi-real photon beams at JLab-12GeV


Coherent tagged Bremsstrahlung:well established technique

→ Hall-B real Bremsstrahlung Photon Tagger



## Performance


- ★  $E_\gamma = 0.8\text{-}5.4 \text{ GeV}$  (20% - 95%  $E_{\text{beam}}$ )
- ★  $\Delta E_\gamma/E_\gamma \sim 10\%$   $t \sim 200\text{ps}$
- ★ Linearly polarized photons  
(coherent Bremsstrahlung)



# Real and quasi-real photon beams at JLab-12GeV


Coherent tagged Bremsstrahlung:well established technique

→ Hall-B real Bremsstrahlung Photon Tagger



## Performance


- ★  $E_\gamma = 0.8\text{-}5.4 \text{ GeV}$  (20% - 95%  $E_{beam}$ )
- ★  $\Delta E_\gamma/E_\gamma \sim 10\%$   $t \sim$
- ★ Linearly polarized (coherent Bremsstrahlung)



Quasi-real electroproduction  
at very Low  $Q^2$






- 
- ★ Test level
  - ★ Fake “ $0^0$ ” electroproduction (no electron in the trigger) from huge collected statistic

Bright meson peaks show up  
The technique works!



# Coherent meson production on nuclei

**Eliminate s-channel resonance background**



**Simplify PWA:** S=I=0 target acts as spin and parity filter for final state mesons

Production cross section expected  $\sim e^{-bt} |A F_A(t)|^2 \rightarrow$  low -t kinematic

**Detection of recoiling nucleus:**


- low -t ( $p \sim 0.2\text{-}0.5$  GeV)
- thin (gas) target ( $\sim 10^{-3}$  g/cm<sup>2</sup>)

**Photon beam:**

- small size
- high flux

**quasi-real photoproduction Hall-B**


## Meson spectroscopy on ${}^4\text{He}$




**Strongest evidence of  $J^{PC}=1^- \pi_1(1400)$  exotic meson**  
 $\pi^- p \rightarrow n \eta \pi^0$  in E852-Brookhaven

**Search for a resonance in P-wave in  $\pi^0 \eta$  and  $\pi^0 \eta'$**


**Known (non-exotic) resonances can be used as a benchmark (e.g.  $J^{PC}=2^{++} a_2(1232)$ )**



# CLAS12 in Hall B





**Existing Hall-B  
tagger**



**Maximum electron angle: 0.5°**  
**The tagger has to be placed upstream to torus supports  
 (option 2)**

**This strongly limits the possible hardware options**

**Two possible options for tagger location:**  
 ★ 1) downstream  
 ★ 2) between target and torus support



# Forward Tagger

## Calorimeter + tracking device

### Electron Energy/momentum

Photon energy ( $\nu = E - E'$ )

Polarization  $\varepsilon^{-1} \sim 1 + \nu^2/2EE'$


PbWO<sub>4</sub> crystals

$R_M \sim 2.2$  cm

$\rho \sim 8.3$  g/cm<sup>3</sup>

$X_0 \sim 0.9$  cm

Low light yield (~1% NaI(Tl))



### Electron angles

$$Q^2 = 4 E E' \sin^2 \vartheta/2$$

$\varphi$  polarization plane

Veto for photons


GEM

Micromegas

SCI-FI hodoscope

Need to estimate resolutions

# CLAS Inner Calorimeter



**424 PbWO<sub>4</sub> crystals**


**L = 16 cm = 17 X<sub>0</sub>**

**Front size 1.3x1.3 cm<sup>2</sup>**

**Back size 1.6x1.6 cm<sup>2</sup>**

**Controlled Temperature (0.1 °C)**


**APD readout**



$$\frac{\sigma_E}{E} = \frac{0.02}{E} \oplus \frac{0.03}{\sqrt{E}} \oplus 0.024$$



$$\sigma_x = \frac{0.2}{\sqrt{E}} \text{ (cm)}$$

# Forward Tagger within CLAS12





## Compatibility with HTCC clearance

- remove HTCC  
no need electron Id
- move HTCC and solenoid upstream (~50cm)  
run parasitically!



# $Q^2$ dependence of the Xsec



Studies at large  $W$  ( $\sim 100 \text{ GeV}$ ) show a smooth transition between  $Q^2 = 0$  and  $Q^2 \neq 0$

## Existing forward taggers

$$Q^2 < W^2$$

COMPASS:	$< 1 \text{ GeV}^2$	$\langle Q^2 \rangle \sim 10^{-1} \text{ GeV}^2$
ZEUS:	$10^{-7} - 0.02 \text{ GeV}^2$	$\langle Q^2 \rangle \sim 5 \cdot 10^{-5} \text{ GeV}^2$
H1:	$< 2 \text{ GeV}^2$	


# Rates in the forward tagger

Inelastic electro-production  
Elastic radiative tail  
Moeller scattering

Signal  
Background

# Rates in the forward tagger

## Inelastic electro-production



## Inelastic electro-production

$\vartheta \ \phi$

Scattered electron angles in the Lab

$v = E_{beam} - E_{e'}$

Quasi-real photon energy

$Q^2 = 4 E E' \sin^2 \vartheta/2$  Virtuality

$\epsilon^{-1} \sim 1 + v^2 / 2 E_{e'} E_{beam}$

Quasi-real photon linear polarization



$$Xsec = \frac{\Gamma_v}{(1+Q^2/.7^2)^2} \sigma_{\gamma p}$$

$$\Gamma_v = \frac{1}{137} \frac{E_{e'}}{2\pi^2 E_{beam}} \frac{W^2 - M_p^2}{2M_p^2} \frac{1}{Q^2} \frac{1}{(1-\epsilon)}$$

$e'$  in the forward tagger


hadrons in CLAS12 ( $N_h >= 1$ )

Rates are limited to  $\sim 10$ kHz




# Rates in the forward tagger

Elastic radiative tail




## Elastic radiative tail



$e'$  in the forward tagger


Elastic proton outside CLAS ( $N_h = 0$  or 1)

Electron rate in the forward tagger is high ( $\sim 1$  MHz)




# Rates in the forward tagger

Moeller scattering




## Moeller scattering

kinematics



Rate




Only 1 electron in the forward tagger

No hits in CLAS ( $N_h = 0$ )

Electron rate in the forward tagger is very high (~50 MHz)

Crystal size:  $0.7 \times 0.7 \text{ cm}^2$

First layer:  
~20x4 crystals



## Hadroproduction kinematic

$L = 130, \Delta\vartheta = 1.9^\circ - 7.0^\circ$

$L_e \sim 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$

$R_e \sim 9.7 \text{ kHz}$

$R_\gamma \sim 0.44 \cdot 10^8 \text{ } \gamma/\text{s}$

$R_e \sim 2.1 \text{ kHz}$

$R_e \sim 1.3 \text{ MHz}$

$R_e \sim 20 \text{ MHz}$

Crystal size =  $1.3 \times 1.3 \text{ cm}^2$

Whole

$N_{\text{crystal}} = 424$

$\Delta\vartheta = 1.9^\circ - 7.0^\circ$

$\Delta E = 7.0 - 10.5 \text{ GeV}$

Hadro production

Eq. photon flux

First layer


$N_{\text{crystal}} = 20$

$\Delta\vartheta(1) = 1.9^\circ - 2.5^\circ$

Hadroproduction

Rad tail ( $\Delta E = 0.3 - 10.9 \text{ GeV}$ )

Moeller



# Conclusions

## Photoproduction experiments at CLAS12

- ★ Many interesting physic topics with a 10 GeV real photon beam
- ★ We are proposing a forward photon tagging facility
- ★ Complementary technique to the Hall-D coherent Bremsstrahlung
- ★ Compatibility with CLAS12 core and concept design identified
- ★ Tagger will be based on PbWO calorimeter + Tracking
- ★ Just started to evaluate rates, background, and hardware designs

We'd like to invite everybody interested in the physics or the hardware design to join us!